



Evidence for a Particle Produced in Association with Weak Bosons and Decaying to a Bottom-Antibottom Quark Pair in Higgs Boson Searches at the Tevatron

T. Aaltonen,^{12,*} V.M. Abazov,^{48,†} B. Abbott,^{112,†} B.S. Acharya,^{31,†} M. Adams,^{78,†} T. Adams,^{74,†} G.D. Alexeev,^{48,†} G. Alkhazov,^{52,†} A. Alton,^{96,†,ff} B. Álvarez González,^{57,*b} G. Alverson,^{92,†} S. Amerio,^{35a,*} D. Amidei,^{96,*} A. Anastassov,^{76,*c} A. Annovi,^{34,*} J. Antos,^{53,*} G. Apollinari,^{76,*} J.A. Appel,^{76,*} T. Arisawa,^{41,*} A. Artikov,^{48,*} J. Asaadi,^{119,*} W. Ashmanskas,^{76,*} A. Askew,^{74,†} S. Atkins,^{89,†} B. Auerbach,^{72,*} K. Augsten,^{9,†} A. Aurisano,^{119,*} C. Avila,^{7,†} F. Afzar,^{66,*} F. Badaud,^{13,†} W. Badgett,^{76,*} T. Bae,^{43,*} L. Bagby,^{76,†} B. Baldin,^{76,†} D.V. Bandurin,^{74,†} S. Banerjee,^{31,†} A. Barbaro-Galtieri,^{68,*} E. Barberis,^{92,†} P. Baringer,^{87,†} V.E. Barnes,^{85,*} B.A. Barnett,^{90,*} P. Barria,^{36c,36a,*} J.F. Bartlett,^{76,†} P. Bartos,^{53,*} U. Bassler,^{18,†} M. Bauce,^{35b,35a,*} V. Bazterra,^{78,†} A. Bean,^{87,†} F. Bedeschi,^{36a,*} M. Begalli,^{2,†} S. Behari,^{90,*} L. Bellantoni,^{76,†} G. Bellettini,^{36b,36a,*} J. Bellinger,^{125,*} D. Benjamin,^{109,*} A. Beretvas,^{76,*} S.B. Beri,^{29,†} G. Bernardi,^{17,†} R. Bernhard,^{22,†} I. Bertram,^{61,†} M. Besançon,^{18,†} R. Beuselinck,^{63,†} P.C. Bhat,^{76,†} S. Bhatia,^{99,†} V. Bhatnagar,^{29,†} A. Bhatti,^{105,*} M. Binkley,^{76,*a} D. Bisello,^{35b,35a,*} I. Bizjak,^{64,*} K.R. Bland,^{122,*} G. Blazey,^{79,†} S. Blessing,^{74,†} K. Bloom,^{100,†} B. Blumenfeld,^{90,*} A. Bocci,^{109,*} A. Bodek,^{106,*} A. Boehnlein,^{76,†} D. Boline,^{107,†} E.E. Boos,^{50,†} G. Borissov,^{61,†} D. Bortoletto,^{85,*} T. Bose,^{91,†} J. Boudreau,^{116,*} A. Boveia,^{77,*} A. Brandt,^{118,†} O. Brandt,^{23,†} L. Brigliadori,^{33b,33a,*} R. Brock,^{98,†} C. Bromberg,^{98,*} A. Bross,^{76,†} D. Brown,^{17,†} J. Brown,^{17,†} E. Brucken,^{12,*} J. Budagov,^{48,*} X.B. Bu,^{76,†} H.S. Budd,^{106,*} M. Buehler,^{76,†} V. Buescher,^{25,†} V. Bunichev,^{50,†} S. Burdin,^{61,†,gg} K. Burkett,^{76,*} G. Busetto,^{35b,35a,*} P. Bussey,^{60,*} C.P. Buszello,^{58,†} A. Buzatu,^{4,*} A. Calamba,^{115,*} C. Calancha,^{56,*} E. Camacho-Pérez,^{45,†} S. Camarda,^{54,†} M. Campanelli,^{64,†} M. Campbell,^{96,*} F. Canelli,^{77,*} B. Carls,^{81,*} D. Carlsmith,^{125,*} R. Carosi,^{36a,*} S. Carrillo,^{73,*d} S. Carron,^{76,*} B. Casal,^{57,*e} M. Casarsa,^{38a,*} B.C.K. Casey,^{76,†} H. Castilla-Valdez,^{45,†} A. Castro,^{33b,33a,*} P. Catastini,^{93,*} S. Caughron,^{98,†} D. Cauz,^{38a,*} V. Cavaliere,^{81,*} M. Cavalli-Sforza,^{54,*} A. Cerri,^{68,*f} L. Cerrito,^{64,*g} S. Chakrabarti,^{107,†} D. Chakraborty,^{79,†} K.M. Chan,^{84,†} A. Chandra,^{121,†} E. Chapon,^{18,†} G. Chen,^{87,†} Y.C. Chen,^{6,*} M. Chertok,^{69,*} S. Chevalier-Théry,^{18,†} G. Chiarelli,^{36a,*} G. Chlachidze,^{76,*} F. Chlebana,^{76,*} D.K. Cho,^{117,†} K. Cho,^{43,*} S.W. Cho,^{44,†} S. Choi,^{44,†} D. Chokheli,^{48,*} B. Choudhary,^{30,†} W.H. Chung,^{125,*} Y.S. Chung,^{106,*} S. Cihangir,^{76,†} M.A. Ciocci,^{36c,36a,*} D. Claes,^{100,†} A. Clark,^{59,*} C. Clarke,^{97,*} J. Clutter,^{87,†} G. Compostella,^{35b,35a,*} M.E. Convery,^{76,*} J. Conway,^{69,*} M. Cooke,^{76,†} W.E. Cooper,^{76,†} M. Corbo,^{76,*} M. Corcoran,^{121,†} M. Cordelli,^{34,*} F. Couderc,^{18,†} M.-C. Cousinou,^{15,†} C.A. Cox,^{69,*} D.J. Cox,^{69,*} F. Crescioli,^{36b,36a,*} A. Croc,^{18,†} J. Cuevas,^{57,*b} R. Culbertson,^{76,*} D. Cutts,^{117,†} D. Dagenhart,^{76,*} N. d'Ascenzo,^{76,*h} A. Das,^{67,†} M. Datta,^{76,*} G. Davies,^{63,†} P. de Barbaro,^{106,*} S.J. de Jong,^{46,47,†} E. De La Cruz-Burelo,^{45,†} F. Déliot,^{18,†} M. Dell'Orso,^{36b,36a,*} R. Demina,^{106,†} L. Demortier,^{105,*} M. Deninno,^{33a,*} D. Denisov,^{76,†} S.P. Denisov,^{51,†} M. d'Errico,^{35b,35a,*} S. Desai,^{76,†} C. Deterre,^{18,†} K. DeVaughan,^{100,†} F. Devoto,^{12,*} A. Di Canto,^{36b,36a,*} B. Di Ruzza,^{76,*} H.T. Diehl,^{76,*} M. Diesburg,^{76,†} P.F. Ding,^{65,†} J.R. Dittmann,^{122,*} A. Dominguez,^{100,†} S. Donati,^{36b,36a,*} P. Dong,^{76,*} M. D'Onofrio,^{62,*} M. Dorigo,^{38a,*} T. Dorigo,^{35a,*} A. Dubey,^{30,†} L.V. Dudko,^{50,†} D. Duggan,^{101,†} A. Duperrin,^{15,†} S. Dutt,^{29,†} A. Dyshkant,^{79,†} M. Eads,^{100,†} K. Ebina,^{41,*} D. Edmunds,^{98,†} A. Elagin,^{119,*} J. Ellison,^{71,†} V.D. Elvira,^{76,†} Y. Enari,^{17,†} A. Eppig,^{96,*} R. Erbacher,^{69,*} S. Errede,^{81,*} N. Ershaidat,^{76,*i} R. Eusebi,^{119,*} H. Evans,^{82,†} A. Evdokimov,^{108,†} V.N. Evdokimov,^{51,†} G. Facini,^{92,†} S. Farrington,^{66,*} M. Feindt,^{24,*} L. Feng,^{79,†} T. Ferbel,^{106,†} J.P. Fernandez,^{56,*} C. Ferrazza,^{37a,*} F. Fiedler,^{25,†} R. Field,^{73,*} F. Filthaut,^{46,47,†} W. Fisher,^{98,†} H.E. Fisk,^{76,†} G. Flanagan,^{76,*j} R. Forrest,^{69,*} M. Fortner,^{79,†} H. Fox,^{61,†} M.J. Frank,^{122,*} M. Franklin,^{93,*} J.C. Freeman,^{76,*} S. Fuess,^{76,†} Y. Funakoshi,^{41,*} M. Gallinaro,^{105,*} A. Garcia-Bellido,^{106,†} J.E. Garcia,^{59,*} J.A. García-González,^{45,†} G.A. García-Guerra,^{45,†,hh} A.F. Garfinkel,^{85,*} P. Garosi,^{36c,36a,*} V. Gavrilov,^{49,†} P. Gay,^{13,†} W. Geng,^{15,98,†} D. Gerbaudo,^{102,†} C.E. Gerber,^{78,†} H. Gerberich,^{81,*} E. Gerchtein,^{76,*} Y. Gershtein,^{101,†} S. Giagu,^{37a,*} V. Giakoumopoulou,^{28,*} P. Giannetti,^{36a,*} K. Gibson,^{116,*} C.M. Ginsburg,^{76,*} G. Ginther,^{76,106,†} N. Giokaris,^{28,*} P. Giromini,^{34,*} G. Giurgiu,^{90,*} V. Glagolev,^{48,*} D. Glenzinski,^{76,*} M. Gold,^{103,*} D. Goldin,^{119,*} N. Goldschmidt,^{73,*} A. Golossanov,^{76,*} G. Golovanov,^{48,†} G. Gomez-Ceballos,^{94,*} G. Gomez,^{57,*} M. Goncharov,^{94,*} O. González,^{56,*} I. Gorelov,^{103,*} A.T. Goshaw,^{109,*} K. Goulianos,^{105,*} A. Goussiou,^{124,†} P.D. Grannis,^{107,†} S. Greder,^{19,†} H. Greenlee,^{76,†} G. Grenier,^{20,†} S. Grinstein,^{54,*} Ph. Gris,^{13,†} J.-F. Grivaz,^{16,†} A. Grohsjean,^{18,†,ii} C. Grosso-Pilcher,^{77,*} R.C. Group,^{123,76,*} S. Grünendahl,^{76,†} M.W. Grünewald,^{32,†} T. Guillemin,^{16,†} J. Guimaraes da Costa,^{93,*} G. Gutierrez,^{76,†} P. Gutierrez,^{112,†} S. Hagopian,^{74,†} S.R. Hahn,^{76,*} J. Haley,^{92,†} E. Halkiadakis,^{101,*} A. Hamaguchi,^{40,*} J.Y. Han,^{106,*} L. Han,^{5,†} F. Happacher,^{34,*} K. Hara,^{42,*} K. Harder,^{65,†} D. Hare,^{101,*} M. Hare,^{95,*} A. Harel,^{106,†} R.F. Harr,^{97,*} K. Hatakeyama,^{122,*} J.M. Hauptman,^{86,†} C. Hays,^{66,*} J. Hays,^{63,†} T. Head,^{65,†} T. Hebbeker,^{21,†} M. Heck,^{24,*} D. Hedin,^{79,†} H. Hegab,^{113,†}

- J. Heinrich,^{114,*} A. P. Heinson,^{71,†} U. Heintz,^{117,†} C. Hensel,^{23,†} I. Heredia-De La Cruz,^{45,†} M. Herndon,^{125,*} K. Herner,^{96,†} G. Hesketh,^{65,†,ji} S. Hewamanage,^{122,*} M. D. Hildreth,^{84,†} R. Hirosky,^{123,†} T. Hoang,^{74,†} J. D. Hobbs,^{107,†} A. Hocker,^{76,*} B. Hoeneisen,^{11,†} J. Hogan,^{121,†} M. Hohlfield,^{25,†} W. Hopkins,^{76,*k} D. Horn,^{24,*} S. Hou,^{6,*} I. Howley,^{118,†} Z. Hubacek,^{9,18,†} R. E. Hughes,^{110,*} M. Hurwitz,^{77,*} U. Husemann,^{72,*} N. Hussain,^{98,*} M. Hussein,^{98,*} J. Huston,^{98,*} V. Hynek,^{9,†} I. Iashvili,^{104,†} Y. Ilchenko,^{120,†} R. Illingworth,^{76,†} G. Introzzi,^{36a,*} M. Iori,^{37b,37a,*} A. S. Ito,^{76,†} A. Ivanov,^{69,*l} S. Jabeen,^{117,†} M. Jaffré,^{16,†} E. James,^{76,*} D. Jang,^{115,*} A. Jayasinghe,^{112,†} B. Jayatilaka,^{109,*} D. T. Jeans,^{37a,*} E. J. Jeon,^{43,*} M. S. Jeong,^{44,†} R. Jesik,^{63,†} P. Jiang,^{5,†} S. Jindariani,^{76,*} K. Johns,^{67,†} E. Johnson,^{98,†} M. Johnson,^{76,†} A. Jonckheere,^{76,†} M. Jones,^{85,*} P. Jonsson,^{63,†} K. K. Joo,^{43,*} J. Joshi,^{71,†} S. Y. Jun,^{115,*} A. W. Jung,^{76,†} T. R. Junk,^{76,*} A. Juste,^{55,†} K. Kaadze,^{88,†} E. Kajfasz,^{15,†} T. Kamon,^{119,*} P. E. Karchin,^{97,*} D. Karmanov,^{50,†} A. Kasmi,^{122,*} P. A. Kasper,^{76,†} Y. Kato,^{40,*m} I. Katsanos,^{100,†} R. Kehoe,^{120,†} S. Kermiche,^{15,†} W. Ketchum,^{77,*} J. Keung,^{114,*} N. Khalatyan,^{76,†} A. Khanov,^{113,†} A. Kharchilava,^{104,†} Y. N. Khazheev,^{48,†} V. Khotilovich,^{119,*} B. Kilminster,^{76,*} D. H. Kim,^{43,*} H. S. Kim,^{43,*} J. E. Kim,^{43,*} M. J. Kim,^{34,*} S. B. Kim,^{43,*} S. H. Kim,^{42,*} Y. J. Kim,^{43,*} Y. K. Kim,^{77,*} N. Kimura,^{41,*} M. Kirby,^{76,*} I. Kiselevich,^{49,†} S. Klimenko,^{73,*} K. Knoepfel,^{76,*} J. M. Kohli,^{29,†} K. Kondo,^{41,*a} D. J. Kong,^{43,*} J. Konigsberg,^{73,*} A. V. Kotwal,^{109,*} A. V. Kozelov,^{51,†} J. Kraus,^{99,†} M. Kreps,^{24,*} J. Kroll,^{114,*} D. Krop,^{77,*} M. Kruse,^{109,*} V. Krutelyov,^{119,*n} T. Kuhr,^{24,*} S. Kulikov,^{51,†} A. Kumar,^{104,†} A. Kupco,^{10,†} M. Kurata,^{42,*} T. Kurča,^{20,†} V. A. Kuzmin,^{50,†} S. Kwang,^{77,*} A. T. Laasanen,^{85,*} S. Lami,^{36a,*} S. Lammel,^{76,*} S. Lammers,^{82,†} M. Lancaster,^{64,*} R. L. Lander,^{69,*} K. Lannon,^{110,*o} A. Lath,^{101,*} G. Latino,^{36c,36a,*} P. Lebrun,^{20,†} T. LeCompte,^{75,*} E. Lee,^{119,*} H. S. Lee,^{44,†} H. S. Lee,^{77,*p} J. S. Lee,^{43,*} S. W. Lee,^{119,*q} S. W. Lee,^{86,†} W. M. Lee,^{76,†} X. Lei,^{67,†} J. Lellouch,^{17,†} S. Leo,^{36b,36a,*} S. Leone,^{36a,*} J. D. Lewis,^{76,*} D. Li,^{17,†} H. Li,^{14,†} L. Li,^{71,†} Q. Z. Li,^{76,†} J. K. Lim,^{44,†} A. Limosani,^{109,*r} D. Lincoln,^{76,†} C.-J. Lin,^{68,*} M. Lindgren,^{76,*} J. Linnemann,^{98,†} V. V. Lipaev,^{51,†} E. Lipeles,^{114,*} R. Lipton,^{76,†} A. Lister,^{59,*} D. O. Litvintsev,^{76,*} C. Liu,^{116,*} H. Liu,^{123,*} H. Liu,^{120,†} Q. Liu,^{85,*} T. Liu,^{76,*} Y. Liu,^{5,†} A. Lobodenko,^{52,†} S. Lockwitz,^{72,*} A. Loginov,^{72,*} M. Lokajicek,^{10,†} R. Lopes de Sa,^{107,†} H. J. Lubatti,^{124,†} D. Lucchesi,^{35b,35a,*} J. Lueck,^{24,*} P. Lujan,^{68,*} P. Lukens,^{76,*} R. Luna-Garcia,^{45,†,kk} G. Lungu,^{105,*} A. L. Lyon,^{76,†} R. Lysak,^{53,*s} J. Lys,^{68,*} A. K. A. Maciel,^{1,*} R. Madar,^{18,†} R. Madrak,^{76,*} P. Maestro,^{36c,36a,*} R. Magaña-Villalba,^{45,†} S. Malik,^{105,*} S. Malik,^{100,†} V. L. Malyshev,^{48,†} G. Manca,^{62,*t} A. Manousakis-Katsikakis,^{28,*} Y. Maravin,^{88,†} F. Margaroli,^{37a,*} C. Marino,^{24,*} M. Martínez,^{54,*} J. Martínez-Ortega,^{45,†} P. Mastrandrea,^{37a,*} K. Matera,^{81,*} M. E. Mattson,^{97,*} A. Mazzacane,^{76,*} P. Mazzanti,^{33a,*} R. McCarthy,^{107,†} K. S. McFarland,^{106,*} C. L. McGivern,^{65,†} P. McIntyre,^{119,*} R. McNulty,^{62,*u} A. Mehta,^{62,*} P. Mehtala,^{12,*} M. M. Meijer,^{46,47,†} A. Melnitchouk,^{99,†} D. Menezes,^{79,†} P. G. Mercadante,^{3,†} M. Merkin,^{50,†} C. Mesropian,^{105,*} A. Meyer,^{21,†} J. Meyer,^{23,†} T. Miao,^{76,*} F. Miconi,^{19,†} D. Mietlicki,^{96,*} A. Mitra,^{6,*} H. Miyake,^{42,*} S. Moed,^{76,*} N. Moggi,^{33a,*} N. K. Mondal,^{31,†} M. N. Mondragon,^{76,*d} C. S. Moon,^{43,*} R. Moore,^{76,*} M. J. Morello,^{36d,36a,*} J. Morlock,^{24,*} P. Movilla Fernandez,^{76,*} A. Mukherjee,^{76,*} M. Mulhearn,^{123,†} Th. Muller,^{24,*} P. Murat,^{76,*} M. Mussini,^{33b,33a,*} J. Nachtman,^{76,*v} Y. Nagai,^{42,*} J. Naganoma,^{41,*} E. Nagy,^{15,†} M. Naimuddin,^{30,†} I. Nakano,^{39,*} A. Napier,^{95,*} M. Narain,^{117,†} R. Nayyar,^{67,†} H. A. Neal,^{96,†} J. P. Negret,^{7,†} J. Nett,^{119,*} M. S. Neubauer,^{81,*} C. Neu,^{123,*} P. Neustroev,^{52,†} H. T. Nguyen,^{123,†} J. Nielsen,^{68,*w} L. Nodulman,^{75,*} S. Y. Noh,^{43,*} O. Norniella,^{81,*} T. Nunnemann,^{26,*} L. Oakes,^{66,*} S. H. Oh,^{109,*} Y. D. Oh,^{43,*} I. Oksuzian,^{123,*} T. Okusawa,^{40,*} R. Orava,^{12,*} J. Orduna,^{121,†} L. Ortolan,^{54,*} N. Osman,^{15,†} J. Osta,^{84,†} M. Padilla,^{71,†} S. Pagan Griso,^{35b,35a,*} C. Pagliarone,^{38a,*} A. Pal,^{118,†} E. Palencia,^{57,*f} V. Papadimitriou,^{76,*} A. A. Paramonov,^{75,*} N. Parashar,^{83,†} V. Parihar,^{117,†} S. K. Park,^{44,†} R. Partridge,^{117,†,ll} N. Parua,^{82,†} J. Patrick,^{76,*} A. Patwa,^{108,†} G. Pauletta,^{38b,38a,*} M. Paulini,^{115,*} C. Paus,^{94,*} D. E. Pellett,^{69,*} B. Penning,^{76,†} A. Penzo,^{38a,*} M. Perfilov,^{50,†} Y. Peters,^{65,†} K. Petridis,^{65,†} G. Petrillo,^{106,†} P. Pétrouff,^{16,†} T. J. Phillips,^{109,*} G. Piacentino,^{36a,*} E. Pianori,^{114,*} J. Pilot,^{110,*} K. Pitts,^{81,*} C. Plager,^{70,*} M.-A. Pleier,^{108,†} P. L. M. Podesta-Lerma,^{45,†,mm} V. M. Podstavkov,^{76,†} L. Pondrom,^{125,*} A. V. Popov,^{51,†} S. Poprocki,^{76,*k} K. Potamianos,^{85,*} A. Pranko,^{68,*} M. Prewitt,^{121,†} D. Price,^{82,†} N. Prokopenko,^{51,†} F. Prokoshin,^{48,*x} F. Ptohos,^{34,*y} G. Punzi,^{36b,36a,*} J. Qian,^{96,†} A. Quadri,^{23,†} B. Quinn,^{99,†} A. Rahaman,^{116,*} V. Ramakrishnan,^{125,*} M. S. Rangel,^{1,†} K. Ranjan,^{30,†} N. Ranjan,^{85,*} P. N. Ratoff,^{61,†} I. Razumov,^{51,†} I. Redondo,^{56,*} P. Renkel,^{120,†} P. Renton,^{66,*} M. Rescigno,^{37a,*} T. Riddick,^{64,*} F. Rimondi,^{33b,33a,*} I. Ripp-Baudot,^{19,†} L. Ristori,^{36a,76,*} F. Rizatdinova,^{113,†} A. Robson,^{60,*} T. Rodriguez,^{114,*} E. Rogers,^{81,*} S. Rolli,^{95,*z} M. Rominsky,^{76,†} R. Roser,^{76,*} A. Ross,^{61,†} C. Royon,^{18,†} P. Rubinov,^{76,†} R. Ruchti,^{84,†} F. Ruffini,^{36c,36a,*} A. Ruiz,^{57,*} J. Russ,^{115,*} V. Rusu,^{76,*} A. Safonov,^{119,*} G. Sajot,^{14,†} W. K. Sakumoto,^{106,*} Y. Sakurai,^{41,*} P. Salcido,^{79,†} A. Sánchez-Hernández,^{45,†} M. P. Sanders,^{26,†} L. Santi,^{38b,38a,*} A. S. Santos,^{1,†,nn} K. Sato,^{42,*} G. Savage,^{76,†} V. Saveliev,^{76,*h} A. Savoy-Navarro,^{76,*aa} L. Sawyer,^{89,†} T. Scanlon,^{63,†} R. D. Schamberger,^{107,†} Y. Scheglov,^{52,†} H. Schellman,^{80,†} P. Schlabach,^{76,*} S. Schlobohm,^{124,*} A. Schmidt,^{24,*} E. E. Schmidt,^{76,*} C. Schwanenberger,^{65,†} T. Schwarz,^{76,*} R. Schwienhorst,^{98,†}

L. Scodellaro,^{57,*} A. Scribano,^{36c,36a,*} F. Scuri,^{36a,*} S. Seidel,^{103,*} Y. Seiya,^{40,*} J. Sekaric,^{87,†} A. Semenov,^{48,*} H. Severini,^{112,†} F. Sforza,^{36b,36a,*} E. Shabalina,^{23,†} S. Z. Shalhout,^{69,*} V. Shary,^{18,†} S. Shaw,^{98,†} A. A. Shchukin,^{51,†} T. Shears,^{62,*} P. F. Shepard,^{116,*} M. Shimojima,^{42,*^{bb}} R. K. Shivpuri,^{30,†} M. Shochet,^{77,*} I. Shreyber-Tecker,^{49,*} V. Simak,^{9,†} A. Simonenko,^{48,*} P. Sinervo,^{4,*} P. Skubic,^{112,†} P. Slattery,^{106,†} K. Sliwa,^{95,*} D. Smirnov,^{84,†} J. R. Smith,^{69,*} K. J. Smith,^{104,†} F. D. Snider,^{76,*} G. R. Snow,^{100,†} J. Snow,^{111,†} S. Snyder,^{108,†} A. Soha,^{76,*} S. Söldner-Rembold,^{65,†} H. Song,^{116,*} L. Sonnenschein,^{21,†} V. Sorin,^{54,*} K. Soustruznik,^{8,†} P. Squillacioti,^{36c,36a,*} R. St. Denis,^{60,*} M. Stancari,^{76,*} J. Stark,^{14,†} O. Stelzer-Chilton,^{4,*} B. Stelzer,^{4,*} D. Stentz,^{76,*^c} D. A. Stoyanova,^{51,†} M. Strauss,^{112,†} J. Strologas,^{103,*} G. L. Strycker,^{96,*} Y. Sudo,^{42,*} A. Sukhanov,^{76,*} I. Suslov,^{48,*} L. Suter,^{65,†} P. Svoisky,^{112,†} M. Takahashi,^{65,†} K. Takemasa,^{42,*} Y. Takeuchi,^{42,*} J. Tang,^{77,*} M. Tecchio,^{96,*} P. K. Teng,^{6,*} J. Thom,^{76,*^k} J. Thome,^{115,*} G. A. Thompson,^{81,*} E. Thomson,^{114,*} M. Titov,^{18,†} D. Toback,^{119,*} S. Tokar,^{53,*} V. V. Tokmenin,^{48,†} K. Tollefson,^{98,*} T. Tomura,^{42,*} D. Tonelli,^{76,*} S. Torre,^{34,*} D. Torretta,^{76,*} P. Totaro,^{35a,*} M. Trovato,^{36d,36a,*} Y.-T. Tsai,^{106,†} K. Tschann-Grimm,^{107,†} D. Tsybychev,^{107,†} B. Tuchming,^{18,†} C. Tully,^{102,†} F. Ukegawa,^{42,*} S. Uozumi,^{43,*} L. Uvarov,^{52,†} S. Uvarov,^{52,†} S. Uzunyan,^{79,†} R. Van Kooten,^{82,†} W. M. van Leeuwen,^{46,†} N. Varelas,^{78,†} A. Varganov,^{96,*} E. W. Varnes,^{67,†} I. A. Vasilyev,^{51,†} F. Vázquez,^{73,*^d} G. Velev,^{76,*} C. Vellidis,^{76,*} P. Verdier,^{20,†} A. Y. Verkheev,^{48,†} L. S. Vertogradov,^{48,†} M. Verzocchi,^{76,†} M. Vesterinen,^{65,†} M. Vidal,^{85,*} I. Vila,^{57,*} D. Vilanova,^{18,†} R. Vilar,^{57,*} J. Vizán,^{57,*} M. Vogel,^{103,*} P. Vokac,^{9,†} G. Volpi,^{34,*} P. Wagner,^{114,*} R. L. Wagner,^{76,*} H. D. Wahl,^{74,†} T. Wakisaka,^{40,*} R. Wallny,^{70,*} S. M. Wang,^{6,*} M. H. L. S. Wang,^{76,†} R.-J. Wang,^{92,†} A. Warburton,^{4,*} J. Warchol,^{84,†} D. Waters,^{64,*} G. Watts,^{124,†} M. Wayne,^{84,†} J. Weichert,^{25,†} L. Welty-Rieger,^{80,†} W. C. Wester III,^{76,*} A. White,^{118,†} D. Whiteson,^{114,*^{cc}} F. Wick,^{24,*} D. Wicke,^{27,†} A. B. Wicklund,^{75,*} E. Wicklund,^{76,*} S. Wilbur,^{77,*} H. H. Williams,^{114,*} M. R. J. Williams,^{61,†} G. W. Wilson,^{87,†} J. S. Wilson,^{110,*} P. Wilson,^{76,*} B. L. Winer,^{110,*} P. Wittich,^{76,*^k} M. Wobisch,^{89,†} S. Wolbers,^{76,*} H. Wolfe,^{110,*} D. R. Wood,^{92,†} T. Wright,^{96,*} X. Wu,^{59,*} Z. Wu,^{122,*} T. R. Wyatt,^{65,†} Y. Xie,^{76,†} R. Yamada,^{76,†} K. Yamamoto,^{40,*} D. Yamato,^{40,*} S. Yang,^{5,†} T. Yang,^{76,*} U. K. Yang,^{77,*^{dd}} W.-C. Yang,^{65,†} Y. C. Yang,^{43,*} W.-M. Yao,^{68,*} T. Yasuda,^{76,†} Y. A. Yatsunenko,^{48,†} W. Ye,^{107,†} Z. Ye,^{76,†} G. P. Yeh,^{76,*} K. Yi,^{76,*^v} H. Yin,^{76,†} K. Yip,^{108,†} J. Yoh,^{76,*} K. Yorita,^{41,*} T. Yoshida,^{40,*^{ee}} S. W. Youn,^{76,†} G. B. Yu,^{109,*} I. Yu,^{43,*} J. M. Yu,^{96,†} S. S. Yu,^{76,*} J. C. Yun,^{76,*} A. Zanetti,^{38a,*} Y. Zeng,^{109,*} J. Zennaro,^{104,†} T. Zhao,^{124,†} T. G. Zhao,^{65,†} B. Zhou,^{96,†} C. Zhou,^{109,*} J. Zhu,^{96,†} M. Zielinski,^{106,†} D. Zieminska,^{82,†} L. Zivkovic,^{117,†} and S. Zucchelli^{33b,33a,*}

(*CDF Collaboration)

(†D0 Collaboration)

¹LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil²Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil³Universidade Federal do ABC, Santo André, Brazil⁴Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3⁵University of Science and Technology of China, Hefei, People's Republic of China⁶Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China⁷Universidad de los Andes, Bogotá, Colombia⁸Charles University, Faculty of Mathematics and Physics, Center for Particle Physics, Prague, Czech Republic⁹Czech Technical University in Prague, Prague, Czech Republic¹⁰Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹¹Universidad San Francisco de Quito, Quito, Ecuador¹²Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France¹⁶LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France¹⁷LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France¹⁸CEA, Ifu, SPP, Saclay, France¹⁹IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany

- ²³*II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany*
- ²⁴*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*
- ²⁵*Institut für Physik, Universität Mainz, Mainz, Germany*
- ²⁶*Ludwig-Maximilians-Universität München, München, Germany*
- ²⁷*Fachbereich Physik, Bergische Universität Wuppertal, Wuppertal, Germany*
- ²⁸*University of Athens, 157 71 Athens, Greece*
- ²⁹*Panjab University, Chandigarh, India*
- ³⁰*Delhi University, Delhi, India*
- ³¹*Tata Institute of Fundamental Research, Mumbai, India*
- ³²*University College Dublin, Dublin, Ireland*
- ^{33a}*Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy*
- ^{33b}*University of Bologna, I-40127 Bologna, Italy*
- ³⁴*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*
- ^{35a}*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- ^{35b}*University of Padova, I-35131 Padova, Italy*
- ^{36a}*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*
- ^{36b}*University of Pisa, I-56127 Pisa, Italy*
- ^{36c}*University of Siena, I-56127 Pisa, Italy*
- ^{36d}*Scuola Normale Superiore, I-56127 Pisa, Italy*
- ^{37a}*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy*
- ^{37b}*Sapienza Università di Roma, I-00185 Roma, Italy*
- ^{38a}*Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, I-33100 Udine, Italy*
- ^{38b}*University of Udine, I-33100 Udine, Italy*
- ³⁹*Okayama University, Okayama 700-8530, Japan*
- ⁴⁰*Osaka City University, Osaka 588, Japan*
- ⁴¹*Waseda University, Tokyo 169, Japan*
- ⁴²*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁴³*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea*
- ⁴⁴*Korea Detector Laboratory, Korea University, Seoul, Korea*
- ⁴⁵*CINVESTAV, Mexico City, Mexico*
- ⁴⁶*Nikhef, Science Park, Amsterdam, the Netherlands*
- ⁴⁷*Radboud University Nijmegen, Nijmegen, the Netherlands*
- ⁴⁸*Joint Institute for Nuclear Research, Dubna, Russia*
- ⁴⁹*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- ⁵⁰*Moscow State University, Moscow, Russia*
- ⁵¹*Institute for High Energy Physics, Protvino, Russia*
- ⁵²*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*
- ⁵³*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*
- ⁵⁴*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*
- ⁵⁵*Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain*
- ⁵⁶*Centro de Investigaciones Energeticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*
- ⁵⁷*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*
- ⁵⁸*Uppsala University, Uppsala, Sweden*
- ⁵⁹*University of Geneva, CH-1211 Geneva 4, Switzerland*
- ⁶⁰*Glasgow University, Glasgow G12 8QQ, United Kingdom*
- ⁶¹*Lancaster University, Lancaster LA1 4YB, United Kingdom*
- ⁶²*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ⁶³*Imperial College London, London SW7 2AZ, United Kingdom*
- ⁶⁴*University College London, London WC1E 6BT, United Kingdom*
- ⁶⁵*The University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁶⁶*University of Oxford, Oxford OX1 3RH, United Kingdom*
- ⁶⁷*University of Arizona, Tucson, Arizona 85721, USA*
- ⁶⁸*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- ⁶⁹*University of California, Davis, Davis, California 95616, USA*
- ⁷⁰*University of California, Los Angeles, Los Angeles, California 90024, USA*

- ⁷¹University of California Riverside, Riverside, California 92521, USA
- ⁷²Yale University, New Haven, Connecticut 06520, USA
- ⁷³University of Florida, Gainesville, Florida 32611, USA
- ⁷⁴Florida State University, Tallahassee, Florida 32306, USA
- ⁷⁵Argonne National Laboratory, Argonne, Illinois 60439, USA
- ⁷⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
- ⁷⁷Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA
- ⁷⁸University of Illinois at Chicago, Chicago, Illinois 60607, USA
- ⁷⁹Northern Illinois University, DeKalb, Illinois 60115, USA
- ⁸⁰Northwestern University, Evanston, Illinois 60208, USA
- ⁸¹University of Illinois, Urbana, Illinois 61801, USA
- ⁸²Indiana University, Bloomington, Indiana 47405, USA
- ⁸³Purdue University Calumet, Hammond, Indiana 46323, USA
- ⁸⁴University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁸⁵Purdue University, West Lafayette, Indiana 47907, USA
- ⁸⁶Iowa State University, Ames, Iowa 50011, USA
- ⁸⁷University of Kansas, Lawrence, Kansas 66045, USA
- ⁸⁸Kansas State University, Manhattan, Kansas 66506, USA
- ⁸⁹Louisiana Tech University, Ruston, Louisiana 71272, USA
- ⁹⁰The Johns Hopkins University, Baltimore, Maryland 21218, USA
- ⁹¹Boston University, Boston, Massachusetts 02215, USA
- ⁹²Northeastern University, Boston, Massachusetts 02115, USA
- ⁹³Harvard University, Cambridge, Massachusetts 02138, USA
- ⁹⁴Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
- ⁹⁵Tufts University, Medford, Massachusetts 02155, USA
- ⁹⁶University of Michigan, Ann Arbor, Michigan 48109, USA
- ⁹⁷Wayne State University, Detroit, Michigan 48201, USA
- ⁹⁸Michigan State University, East Lansing, Michigan 48824, USA
- ⁹⁹University of Mississippi, University, Mississippi 38677, USA
- ¹⁰⁰University of Nebraska, Lincoln, Nebraska 68588, USA
- ¹⁰¹Rutgers University, Piscataway, New Jersey 08855, USA
- ¹⁰²Princeton University, Princeton, New Jersey 08544, USA
- ¹⁰³University of New Mexico, Albuquerque, New Mexico 87131, USA
- ¹⁰⁴State University of New York, Buffalo, New York 14260, USA
- ¹⁰⁵The Rockefeller University, New York, New York 10065, USA
- ¹⁰⁶University of Rochester, Rochester, New York 14627, USA
- ¹⁰⁷State University of New York, Stony Brook, New York 11794, USA
- ¹⁰⁸Brookhaven National Laboratory, Upton, New York 11973, USA
- ¹⁰⁹Duke University, Durham, North Carolina 27708, USA
- ¹¹⁰The Ohio State University, Columbus, Ohio 43210, USA
- ¹¹¹Langston University, Langston, Oklahoma 73050, USA
- ¹¹²University of Oklahoma, Norman, Oklahoma 73019, USA
- ¹¹³Oklahoma State University, Stillwater, Oklahoma 74078, USA
- ¹¹⁴University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ¹¹⁵Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA
- ¹¹⁶University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
- ¹¹⁷Brown University, Providence, Rhode Island 02912, USA
- ¹¹⁸University of Texas, Arlington, Texas 76019, USA
- ¹¹⁹Texas A&M University, College Station, Texas 77843, USA
- ¹²⁰Southern Methodist University, Dallas, Texas 75275, USA
- ¹²¹Rice University, Houston, Texas 77005, USA
- ¹²²Baylor University, Waco, Texas 76798, USA
- ¹²³University of Virginia, Charlottesville, Virginia 22904, USA
- ¹²⁴University of Washington, Seattle, Washington 98195, USA
- ¹²⁵University of Wisconsin, Madison, Wisconsin 53706, USA

(Received 26 July 2012; published 14 August 2012)

We combine searches by the CDF and D0 Collaborations for the associated production of a Higgs boson with a W or Z boson and subsequent decay of the Higgs boson to a bottom-antibottom quark pair. The data, originating from Fermilab Tevatron $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, correspond to integrated luminosities of up to 9.7 fb^{-1} . The searches are conducted for a Higgs boson with mass in the range

100–150 GeV/ c^2 . We observe an excess of events in the data compared with the background predictions, which is most significant in the mass range between 120 and 135 GeV/ c^2 . The largest local significance is 3.3 standard deviations, corresponding to a global significance of 3.1 standard deviations. We interpret this as evidence for the presence of a new particle consistent with the standard model Higgs boson, which is produced in association with a weak vector boson and decays to a bottom-antibottom quark pair.

DOI: [10.1103/PhysRevLett.109.071804](https://doi.org/10.1103/PhysRevLett.109.071804)

PACS numbers: 13.85.Rm, 14.80.Bn

The standard model (SM) [1,2] Higgs boson H is predicted to be produced in association with a W or Z boson at the Fermilab Tevatron $p\bar{p}$ Collider if it is within kinematic reach, and its dominant decay mode is predicted to be into a bottom-antibottom quark pair ($b\bar{b}$), if its mass m_H is less than 135 GeV/ c^2 [3,4]. An observation of this process would support the SM prediction that the mechanism for electroweak symmetry breaking, which gives mass to the weak vector bosons, is also the source of fermionic mass in the quark sector. The leptonic decays of the W and Z vector bosons and the decays of the H to $b\bar{b}$ provide distinctive signatures of Higgs boson production, which are used to discriminate signal events from the copious backgrounds [5]. In this Letter, we combine the searches from the CDF and D0 Collaborations for H bosons produced in association with a vector boson, with subsequent decays $H \rightarrow b\bar{b}$. Both collaborations consider the processes $WH \rightarrow \ell\nu b\bar{b}$, $ZH \rightarrow \ell^+\ell^- b\bar{b}$, and $WH, ZH \rightarrow \cancel{E}_T b\bar{b}$ [6–11] (where ℓ is either e or μ and \cancel{E}_T denotes missing transverse energy [12]), and separately combine results within their collaborations [13,14]. This is the first publication of a combination of CDF and D0's searches for $H \rightarrow b\bar{b}$, which is based on the preliminary findings reported in Ref. [15].

Much is known about the Higgs boson from other experiments. The direct searches at LEP2 in the $e^+e^- \rightarrow ZH(\rightarrow b\bar{b})$ mode, with a small contribution from vector boson fusion, are very similar to those combined here, and exclude SM Higgs boson masses below 114.4 GeV/ c^2 at the 95% confidence level (C.L.) [16]. Direct searches for $VH \rightarrow Vb\bar{b}$ at the LHC, where $V = W$ or Z [17,18], do not yet constrain the allowed SM Higgs boson mass range. Including other search modes, direct searches at the LHC for the SM Higgs boson limit its mass to be between 116.6 and 119.4 GeV/ c^2 or between 122.1 and 127.0 GeV/ c^2 , at the 95% C.L. [19,20]. Within these searches, both LHC experiments observe local excesses above the background expectations for a Higgs boson mass of approximately 125 GeV/ c^2 . With additional data and analysis improvements, the LHC experiments confirm these excesses and observe a particle with properties consistent with those predicted for the SM Higgs boson [21]. Much of the power of the LHC searches comes from $gg \rightarrow H$ production and Higgs boson decays to $\gamma\gamma$, W^+W^- , and ZZ , which probe the couplings of the Higgs boson to other bosons. In the allowed mass range, the Tevatron experiments are particularly sensitive to VH production with $H \rightarrow b\bar{b}$, which

probes the Higgs boson's coupling to b quarks. We search for Higgs bosons of masses $100 < m_H < 150$ GeV/ c^2 and interpret our results independently of searches which are not sensitive to the specific Higgs boson production and decay modes studied here. We also report results assuming $m_H = 125$ GeV/ c^2 .

Higgs boson signal events are simulated using the leading order (LO) calculation from PYTHIA [22], with CTEQ5L (CDF) and CTEQ6L1 (D0) [23] parton distribution functions (PDFs). We normalize our Higgs boson signal-rate predictions to the highest-order calculations available. The WH and ZH cross section calculations are performed at next-to-next-to leading order (NNLO) precision in QCD and next-to-leading-order (NLO) precision in the electroweak corrections [3]. We use the branching fractions for Higgs boson decay from Ref. [4]. These rely on calculations using HDECAY [24] and PROPHECY4F [25]. Assuming the $m_H = 125$ GeV/ c^2 hypothesis, we expect approximately 155 Higgs boson signal events to pass our selection requirements, along with 9.2×10^4 background events from all other SM sources.

We model SM and instrumental background processes using a mixture of Monte Carlo (MC) and data-driven methods. For CDF, backgrounds from SM processes with electroweak gauge bosons or top quarks are modeled using PYTHIA, ALPGEN [26], MC@NLO [27], and HERWIG [28]. For D0, these backgrounds are modeled using PYTHIA, ALPGEN, and COMPEP [29]. An interface to PYTHIA provides parton showering and hadronization for generators without this functionality.

Diboson (WW , WZ , ZZ) MC samples are normalized using the NLO calculations from MCFM [30]. For $t\bar{t}$, we use a production cross section of 7.04 ± 0.70 pb [31], which is based on a top-quark mass of 173 GeV/ c^2 [32] and MSTW 2008 NNLO PDFs [33]. The single-top-quark production cross section is taken to be 3.15 ± 0.31 pb [34]. Data-driven methods are used to normalize the W/Z plus light-flavor and heavy-flavor jet backgrounds [35] using W/Z data events containing no b -tagged jets [36], which have negligible signal content [13,14].

The CDF and D0 detectors are multipurpose solenoidal spectrometers surrounded by hermetic calorimeters and muon detectors and are designed to study the products of 1.96 TeV proton-antiproton collisions [37,38]. All searches combined here use the complete Tevatron data sample, which after data quality requirements corresponds to

9.45 fb^{-1} – 9.7 fb^{-1} ; the size of the analyzed data set depends on the experiment and the search channel. The online event selections (triggers) rely on fast reconstruction of combinations of high- p_T lepton candidates, jets, and \cancel{E}_T . Event selections are similar in the CDF and D0 analyses, consisting typically of a preselection based on event topology and kinematics, and a subsequent selection using b -tagging. Both collaborations use multivariate analysis (MVA) techniques that combine several discriminating variables into a single final discriminant which is used to separate signal from background. Each channel is divided into exclusive subchannels according to various lepton, jet multiplicity, and b -tagging characterization criteria aimed at grouping events with similar signal-to-background ratio and so optimize the overall sensitivity. Because of the importance of b -tagging, both collaborations have developed multivariate approaches to maximize the performance of the b -tagging algorithms. A boosted decision tree algorithm is used in the D0 analysis, which builds and improves upon the previous neural network b -tagger [39], giving an identification efficiency of $\approx 80\%$ for b -jets with a misidentification rate of $\approx 10\%$. The CDF b -tagging algorithm has been recently augmented with an MVA [40], providing a b -tagging efficiency of $\approx 70\%$ and a misidentification rate of $\approx 5\%$.

The reconstructed dijet mass provides discrimination between signal and background. The decay width of the Higgs boson is expected to be much narrower than the experimental dijet mass resolution, which is typically 15% of the mean reconstructed mass. A SM Higgs boson signal would appear as a broad enhancement in the reconstructed dijet mass distribution. The sensitivity is enhanced by combining the dijet mass with other kinematic information using multivariate discriminants. The MVA functions are optimized separately for each subchannel and for each hypothesized value of m_H in the range 100–150 GeV/ c^2 , in 5 GeV/ c^2 intervals. The results from each subchannel are summarized in histograms of the MVA discriminants for the expected Higgs boson signals, the backgrounds itemized by source, and the observed data.

We interpret the results using both Bayesian and modified frequentist techniques, separately at each value of m_H . These methods are described in Refs. [15,41,42]. These techniques are built on a likelihood function which is a product of Poisson probabilities for observing the data in each bin of each subchannel. Systematic uncertainties are parametrized with nuisance parameters, which affect the rates of the predicted signal and background yields in each bin. A nuisance parameter may affect the predictions of multiple sources of signal and background in multiple subchannels, thus taking correlations into account. A nuisance parameter may also affect multiple bins' predictions by different amounts, thus parameterizing uncertainty in the shapes of distributions. Gaussian priors are assumed for the nuisance parameters, truncated to ensure that no pre-

diction is negative. The signal predictions used correspond to SM Higgs boson production and decay, scaled by a factor R for all bins of all subchannels. By scaling all signal contributions by the same factor, we assume that the relative contributions of the different processes are as predicted by the SM.

In the Bayesian technique, we assume a uniform prior in R and integrate the likelihood function multiplied by the priors of the nuisance parameters to obtain the posterior density for R . The observed 95% credibility level upper limit on R , R_{95}^{obs} , is such that 95% of the integral of the posterior of R is below R_{95}^{obs} . The expected distribution of R_{95} is computed in an ensemble of simulated experimental outcomes assuming no signal is present. In each simulated outcome, random values of the nuisance parameters are drawn from their priors. A combined measurement of the cross section for Higgs boson production times the branching fraction $\mathcal{B}(H \rightarrow b\bar{b})$, in units of the SM production rate, is given by R^{fit} , which is the value of R that maximizes the posterior density. The 68% credibility interval, which corresponds to 1 standard deviation (s.d.), is quoted as the smallest interval containing 68% of the integral of the posterior.

We also perform calculations using the modified frequentist technique [42], CL_s , using a log-likelihood ratio (LLR) as the test statistic:

$$LLR = -2 \ln \frac{p(\text{data}|H_1)}{p(\text{data}|H_0)}, \quad (1)$$

where H_1 denotes the test hypothesis, which admits the presence of SM backgrounds and a Higgs boson signal, H_0 denotes the null hypothesis, for only SM backgrounds, and “data” are either simulated data constructed from the expected signal and backgrounds, or the actual observed data. The probabilities p are computed using the best-fit posterior values of the nuisance parameters for each simulated experimental outcome, separately for each of the two hypotheses, and include the Poisson probabilities of observing the data multiplied by Gaussian constraint terms for the values of the nuisance parameters. The CL_s technique involves computing two p values,

$$\text{CL}_b = p(LLR \geq LLR_{\text{obs}}|H_0), \quad (2)$$

where LLR_{obs} is the value of the test statistic computed for the data, and

$$\text{CL}_{s+b} = p(LLR \geq LLR_{\text{obs}}|H_1). \quad (3)$$

To compute limits, we use the ratio of p values, $\text{CL}_s = \text{CL}_{s+b}/\text{CL}_b$. If $\text{CL}_s < 0.05$ for a particular choice of H_1 , parametrized by the signal scale factor R , that hypothesis is excluded at the 95% C. L. The median expected limit is computed using the median LLR value expected in the background-only hypothesis.

The uncertainties on the signal production cross sections are estimated from the factorization and renormalization scale variations, which include the impact of uncalculated

higher-order corrections, as well as uncertainties due to PDFs, and the dependence on the strong coupling constant, α_s . The resulting uncertainties on the inclusive WH and ZH production rates are 7% [3]. We assign uncertainties to the prediction of $\mathcal{B}(H \rightarrow b\bar{b})$ as calculated in Ref. [43]. These uncertainties arise from imperfect knowledge of the mass of the b and c quarks, α_s , and theoretical uncertainties in the $b\bar{b}$ and W^+W^- decay rates.

The largest sources of uncertainty on the dominant backgrounds are the rates of tagged V + heavy flavor jets, which are typically 20%–30% of the predicted values. The posterior uncertainties on these rates are typically 8% or less. Uncertainties on lepton identification and trigger efficiencies range from 2% to 6% and are applied to both signal- and MC-based background predictions. These uncertainties are estimated from data-based methods separately by CDF and D0, and differ based on lepton flavor and identification category. The b -tag efficiencies and mistag rates are similarly constrained by auxiliary data samples, such as inclusive jet data or $t\bar{t}$ events. The uncertainty on the per-jet b -tag efficiency is approximately 4%, and the mistag uncertainties vary between 7% and 15%. The uncertainties on the measurements of the integrated luminosities, which are used to normalize the expected signal yields and the MC-based backgrounds, are 6% (CDF) [44] and 6.1% (D0) [45]. Of these values, 4% arises from the inelastic $p\bar{p}$ cross section, which is taken to be correlated between CDF and D0.

To validate our background modeling and search methods, we perform a search for SM diboson production in the same final states used for the SM $H \rightarrow b\bar{b}$ searches. The NLO SM cross section for VZ times the branching fraction of $Z \rightarrow b\bar{b}$ is 0.68 ± 0.05 pb, which is about 6 times larger than the 0.12 ± 0.01 pb cross section times branching fraction of $VH(H \rightarrow b\bar{b})$ for a $125 \text{ GeV}/c^2$ SM Higgs boson. The data sample, reconstruction, process modeling, uncertainties, and subchannel divisions are identical to those of the SM Higgs boson search. However, discriminant functions are trained to distinguish the contributions of SM diboson production from those of other backgrounds, and potential contributions from Higgs boson production are not considered. The measured cross section for VZ is $3.9 \pm 0.6(\text{stat}) \pm 0.7(\text{syst})$ pb, which is consistent with the SM prediction of 4.4 ± 0.3 pb.

The combined background-subtracted reconstructed dijet mass (m_{jj}) distribution for the VZ analysis is shown in Fig. 1. The VZ signal and the background contributions are fit to the data, and the fitted background is subtracted. Also shown is the contribution expected from a SM Higgs boson with $m_H = 125 \text{ GeV}/c^2$.

To visualize the results produced by the multivariate VH analyses, we combine the histograms of the final discriminants, adding the contents of bins with similar signal-to-background ratio (s/b). Figure 2 shows the signal expectation and the data with the background (including




FIG. 1 (color online). Background-subtracted distribution of the reconstructed dijet mass m_{jj} , summed over all input channels. The VZ signal and the background contributions are fit to the data, and the fitted background is subtracted. The fitted VZ and expected SM Higgs ($m_H = 125 \text{ GeV}/c^2$) contributions are shown with filled histograms.

VZ) subtracted, as a function of the s/b of the collected bins, for the combined Higgs boson search, assuming $m_H = 125 \text{ GeV}/c^2$. The background model is fit to the data, and the uncertainties on the background are those after the nuisance parameters have been constrained in the fit. An excess of events in the highest s/b bins relative to the background-only expectation is observed. We also show the LLR as a function of m_H in Fig. 3, along with its expected values under the hypotheses H_0 and H_1 , and also the hypothesis that a SM Higgs boson is present with $m_H = 125 \text{ GeV}/c^2$.

We extract limits on SM Higgs boson production as a function of m_H in the range 100–150 GeV/c^2 in terms of R_{95}^{obs} , the observed limit relative to the SM rate. These




FIG. 2 (color online). Background-subtracted distribution for the discriminant histograms, summed for bins with similar signal-to-background ratio (s/b), for the $H \rightarrow b\bar{b}$ ($m_H = 125 \text{ GeV}/c^2$) search. The solid histogram shows the uncertainty on the background after the fit to the data as discussed in the text. The signal model, scaled to the SM expectation, is shown with a filled histogram. Uncertainties on the data points correspond to the square root of the sum of the expected signal and background yields in each bin.




FIG. 3 (color online). The log-likelihood ratio LLR as a function of Higgs boson mass. The dark and light-shaded bands correspond to the regions encompassing 1 s.d. and 2 s.d. fluctuations of the background, respectively. The dot-dashed line shows the median expected LLR assuming the SM Higgs boson is present at $m_H = 125 \text{ GeV}/c^2$.

limits are shown in Fig. 4, together with the median expected values and distributions in simulated experimental outcomes assuming a signal is absent. We also show the median expected limits assuming the SM Higgs boson, with $m_H = 125 \text{ GeV}/c^2$, is present. We exclude $m_H < 106 \text{ GeV}/c^2$ at the 95% credibility level, while our median expected limit on m_H is $116 \text{ GeV}/c^2$, if no signal were present. The exclusions obtained with the CL_s technique match those computed with the Bayesian technique.

The observed limits are weaker than expected due to an excess events in the data with respect to the background predictions in the most sensitive bins of the discriminant distributions, favoring the hypothesis that a signal is present. We characterize this excess by computing the best-fit rate parameter R^{fit} , which, when multiplied by the SM prediction for the associated production cross section times




FIG. 4 (color online). The observed 95% credibility level upper limits on SM Higgs boson production (R_{95}) as a function of Higgs boson mass. The dashed line indicates the median expected value in the absence of a signal, and the shaded bands indicate the 1 s.d. and 2 s.d. ranges in which R_{95} is expected to fluctuate. The dot-dashed line shows the median expected limit if the SM Higgs boson is present at $m_H = 125 \text{ GeV}/c^2$.

the decay branching ratio $(\sigma_{WH} + \sigma_{ZH}) \times \mathcal{B}(H \rightarrow b\bar{b})$, yields the best-fit value for this quantity. We show our fitted $(\sigma_{WH} + \sigma_{ZH}) \times \mathcal{B}(H \rightarrow b\bar{b})$ as a function of m_H , along with the SM prediction, in Fig. 5. The figure also shows the expected cross section fits for each m_H assuming that the SM Higgs boson, with $m_H = 125 \text{ GeV}/c^2$, is present. The expected fits are shown for both the expected SM rate and the best fitted rate from data, which corresponds to $(\sigma_{WH} + \sigma_{ZH}) \times \mathcal{B}(H \rightarrow b\bar{b}) = 0.23^{+0.09}_{-0.08} \text{ (stat + syst) pb}$. The corresponding SM prediction for $m_H = 125 \text{ GeV}/c^2$ is $0.12 \pm 0.01 \text{ pb}$.

The significance of the excess in the data over the background prediction is computed at each hypothesized Higgs boson mass in the range 100–150 GeV/c^2 by calculating the local p value under the background-only hypothesis using R^{fit} as the test statistic. This p value expresses the probability to obtain the value of R^{fit} observed in the data or larger, assuming a signal is truly absent. These p -values are shown in Fig. 6 along with the expected p values assuming a SM signal is present, separately for each value of m_H . The observed p value as a function of m_H exhibits a broad minimum and the maximum local significance corresponds to 3.3 standard deviations at $m_H = 135 \text{ GeV}/c^2$.

The Look-Elsewhere Effect (LEE) [46,47] accounts for the possibility of a background fluctuation affecting the local p -value anywhere in the tested m_H range. In the mass range from 115 GeV/c^2 (the prior bound from the LEP2 direct search [16]) to 150 GeV/c^2 , the reconstructed mass resolution is typically 15%, and the resulting LEE factor is approximately 2. Correcting for the LEE yields a global significance of 3.1 standard deviations. Taking into account




FIG. 5 (color online). The best-fit cross section times branching ratio $(\sigma_{WH} + \sigma_{ZH}) \times \mathcal{B}(H \rightarrow b\bar{b})$ as a function of m_H . The dark and light-shaded regions indicate the 1 s.d. and 2 s.d. measurement uncertainties, and the SM prediction is shown as the smooth, falling curve with a narrow band indicating the theoretical uncertainty. The expected cross section fit values assuming the SM Higgs boson is present at $m_H = 125 \text{ GeV}/c^2$ are shown with dot-dashed lines for the cases of the expected SM rate (dark blue) and the best fitted rate from data (light magenta).




FIG. 6 (color online). The p value as a function of m_H under the background-only hypothesis. Also shown are the median expected values assuming a SM signal is present, evaluated separately at each m_H . The associated dark and light-shaded bands indicate the 1 s.d. and 2 s.d. fluctuations of possible experimental outcomes.

the exclusion limits for the SM Higgs boson mentioned earlier, there is no LEE and we derive a significance of 2.8 standard deviations for $m_H = 125 \text{ GeV}/c^2$.

We interpret this result as evidence for the presence of a particle that is produced in association with a W or Z boson and decays to a bottom-antibottom quark pair. The excess seen in the data is most significant in the mass range between 120 and 135 GeV/c^2 , and is consistent with production of the SM Higgs boson within this mass range. Assuming a Higgs boson exists in this mass range, these results provide a direct probe of its coupling to b quarks.

We thank the Fermilab staff and technical staffs of the participating institutions for their vital contributions and acknowledge support from the DOE and NSF (USA), ARC (Australia), CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil), NSERC (Canada), NSC, CAS and CNSF (China), Colciencias (Colombia), MSMT and GACR (Czech Republic), the Academy of Finland, CEA and CNRS/IN2P3 (France), BMBF and DFG (Germany), DAE and DST (India), SFI (Ireland), INFN (Italy), MEXT (Japan), the Korean World Class University Program and NRF (Korea), CONACyT (Mexico), FOM (Netherlands), MON, NRC KI and RFBR (Russia), the Slovak R&D Agency, the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010 (Spain), The Swedish Research Council (Sweden), SNSF (Switzerland), STFC and the Royal Society (United Kingdom), and the A. P Sloan Foundation (USA).

^aDeceased.

^bCDF visitor from Universidad de Oviedo, E-33007 Oviedo, Spain.

^cCDF visitor from Northwestern University, Evanston, IL 60208, USA.

^dCDF visitor from Universidad Iberoamericana, Mexico D.F., Mexico.

^eCDF visitor from ETH, 8092 Zurich, Switzerland.

^fCDF visitor from CERN, CH-1211 Geneva, Switzerland.

^gCDF visitor from Queen Mary, University of London, London, E1 4NS, United Kingdom.

^hCDF visitor from National Research Nuclear University, Moscow, Russia.

ⁱCDF visitor from Yarmouk University, Irbid 211-63, Jordan.

^jCDF visitor from Muons, Inc., Batavia, IL 60510, USA.

^kCDF visitor from Cornell University, Ithaca, NY 14853, USA.

^lCDF visitor from Kansas State University, Manhattan, KS 66506, USA.

^mCDF visitor from Kinki University, Higashi-Osaka City, Japan 577-8502.

ⁿCDF visitor from University of California Santa Barbara, Santa Barbara, CA 93106, USA.

^oCDF visitor from University of Notre Dame, Notre Dame, IN 46556, USA.

^pCDF visitor from Ewha Womans University, Seoul, 120-750, Korea.

^qCDF visitor from Texas Tech University, Lubbock, TX 79609, USA.

^rCDF visitor from University of Melbourne, Victoria 3010, Australia.

^sCDF visitor from Institute of Physics, Academy of Sciences of the Czech Republic, Czech Republic.

^tCDF visitor from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy.

^uCDF visitor from University College Dublin, Dublin 4, Ireland.

^vCDF visitor from University of Iowa, Iowa City, IA 52242, USA.

^wCDF visitor from University of California, Santa Cruz, Santa Cruz, CA 95064, USA.

^xCDF visitor from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile.

^yCDF visitor from University of Cyprus, Nicosia CY-1678, Cyprus.

^zCDF visitor from Office of Science, U.S. Department of Energy, Washington, DC 20585, USA.

^{aa}CDF visitor from CNRS-IN2P3, Paris, F-75205 France.

^{bb}CDF visitor from Nagasaki Institute of Applied Science, Nagasaki, Japan.

^{cc}CDF visitor from University of California, Irvine, Irvine, CA 92697, USA.

^{dd}CDF visitor from University of Manchester, Manchester M13 9PL, United Kingdom.

^{ee}CDF visitor from University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017.

^{ff}D0 visitor from Augustana College, Sioux Falls, SD, USA.

^{gg}D0 visitor from The University of Liverpool, Liverpool, UK.

^{hh}D0 visitor from UPIITA-IPN, Mexico City, Mexico.

- ⁱⁱD0 visitor from DESY, Hamburg, Germany.
- ^{jj}D0 visitor from University College London, London, UK.
- ^{kk}D0 visitor from Centro de Investigacion en Computacion - IPN, Mexico City, Mexico.
- ^{ll}D0 visitor from SLAC, Menlo Park, CA, USA.
- ^{mmm}D0 visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.
- ⁿⁿD0 visitor from Universidade Estadual Paulista, São Paulo, Brazil.
- [1] S.L. Glashow, *Nucl. Phys.* **22**, 579 (1961); S. Weinberg, *Phys. Rev. Lett.* **19**, 1264 (1967); A. Salam, *Elementary Particle Theory*, edited by N. Svartholm (Almqvist and Wiksell, Stockholm 1968), p. 367.
 - [2] F. Englert and R. Brout, *Phys. Rev. Lett.* **13**, 321 (1964); P.W. Higgs, *Phys. Rev. Lett.* **13**, 508 (1964); G.S. Guralnik, C.R. Hagen, and T.W.B. Kibble, *Phys. Rev. Lett.* **13**, 585 (1964); P.W. Higgs, *Phys. Rev.* **145**, 1156 (1966).
 - [3] J. Baglio and A. Djouadi, *J. High Energy Phys.* **10** (2010) 064; O. Brein, R.V. Harlander, M. Weisemann, and T. Zirke, *Eur. Phys. J. C* **72**, 1868 (2012).
 - [4] S. Dittmaier *et al.* (LHC Higgs Cross Section Working Group), [arXiv:1201.3084](#).
 - [5] A. Stange, W. Marciano, and S. Willenbrock, *Phys. Rev. D* **49**, 1354 (1994); A. Stange, W. Marciano, and S. Willenbrock, *Phys. Rev. D* **50**, 4491 (1994).
 - [6] T. Aaltonen *et al.* (CDF Collaboration), [arXiv:1207.1703](#) [Phys. Rev. Lett. (to be published)].
 - [7] T. Aaltonen *et al.* (CDF Collaboration), [arXiv:1207.1711](#) [Phys. Rev. Lett. (to be published)].
 - [8] T. Aaltonen *et al.* (CDF Collaboration), [arXiv:1207.1704](#) [Phys. Rev. Lett. (to be published)].
 - [9] V.M. Abazov *et al.* (D0 Collaboration), [arXiv:1208.0653](#) [Phys. Rev. Lett. (to be published)].
 - [10] V.M. Abazov *et al.* (D0 Collaboration), [arXiv:1207.5689](#) [Phys. Lett. B (to be published)].
 - [11] V.M. Abazov *et al.* (D0 Collaboration), [arXiv:1207.5819](#) [Phys. Rev. Lett. (to be published)].
 - [12] CDF and D0 use cylindrical coordinate systems with origins in the centers of the detectors, where θ and ϕ are the polar and azimuthal angles, respectively, and pseudorapidity is $\eta = -\ln \tan(\theta/2)$. The transverse energy, as measured by the calorimetry, is defined to be $E_T = E \sin\theta$. The missing E_T (\cancel{E}_T) is defined by $\cancel{E}_T = -\sum_i E_i^j \hat{n}_i$, i = calorimeter tower number, where \hat{n}_i is a unit vector perpendicular to the beam axis and pointing at the i th calorimeter tower. \cancel{E}_T is corrected for high-energy muons and also jet energy corrections. We define $\cancel{E}_T = |\cancel{E}_T|$. The transverse momentum p_T is defined to be $p \sin\theta$.
 - [13] T. Aaltonen *et al.* (CDF Collaboration), [arXiv:1207.1707](#) [Phys. Rev. Lett. (to be published)].
 - [14] V.M. Abazov *et al.* (D0 Collaboration), [arXiv:1207.6631](#) [Phys. Rev. Lett. (to be published)].
 - [15] The CDF and D0 Collaborations and the Tevatron New Physics and Higgs Working Group, [arXiv:1207.0449](#).
 - [16] The ALEPH, DELPHI, L3 and OPAL Collaborations, and the LEP Working Group for Higgs Boson Searches, *Phys. Lett. B* **565**, 61 (2003).
 - [17] G. Aad *et al.* (ATLAS Collaboration), [arXiv:1207.0210](#) [Phys. Lett. B (to be published)].
 - [18] S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Lett. B* **710**, 284 (2012).
 - [19] S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Lett. B* **710**, 26 (2012).
 - [20] G. Aad *et al.* (ATLAS Collaboration), [arXiv:1207.0319](#) [Phys. Rev. D (to be published)].
 - [21] G. Aad *et al.* (ATLAS Collaboration), [arXiv:1207.7214](#) [Phys. Lett. B (to be published)]; S. Chatrchyan *et al.* (CMS Collaboration), [arXiv:1207.7235](#) [Phys. Lett. B (to be published)].
 - [22] T. Sjöstrand, S. Mrenna, and P. Skands, *J. High Energy Phys.* **05** (2006) 026. We use PYTHIA version 6.216 to generate the Higgs boson signals.
 - [23] H.L. Lai *et al.*, *Eur. Phys. J. C* **12**, 375 (2000); J. Pumplin *et al.*, *J. High Energy Phys.* **07** (2002) 012.
 - [24] A. Djouadi, J. Kalinowski, and M. Spira, *Comput. Phys. Commun.* **108**, 56 (1998).
 - [25] A. Bredenstein, A. Denner, S. Dittmaier, and M.M. Weber, *Phys. Rev. D* **74**, 013004 (2006); A. Bredenstein, A. Denner, S. Dittmaier, A. Mück, and M.M. Weber, *J. High Energy Phys.* **02** (2007) 080.
 - [26] M. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. Polosa, *J. High Energy Phys.* **07** (2003) 001.
 - [27] S. Frixione and B.R. Webber, *J. High Energy Phys.* **06** (2002) 029.
 - [28] G. Corcella *et al.*, *J. High Energy Phys.* **01** (2001) 010.
 - [29] A. Pukhov *et al.*, [arXiv:hep-ph/9908288](#); E. Boos *et al.* (CompHEP Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **534**, 250 (2004); E. Boos, V.E. Bunichev, L.V. Dudko, V.I. Savrin, and A.V. Sherstnev, *Phys. At. Nucl.* **69**, 1317 (2006).
 - [30] J.M. Campbell and R.K. Ellis, *Phys. Rev. D* **60**, 113006 (1999).
 - [31] U. Langenfeld, S. Moch, and P. Uwer, *Phys. Rev. D* **80**, 054009 (2009).
 - [32] T. Aaltonen *et al.* (CDF and D0 Collaborations), [arXiv:1207.1069](#) [Phys. Rev. D (to be published)].
 - [33] A.D. Martin, W.J. Stirling, R.S. Thorne, and G. Watt, *Eur. Phys. J. C* **63**, 189 (2009).
 - [34] N. Kidonakis, *Phys. Rev. D* **74**, 114012 (2006).
 - [35] A heavy-flavor jet is a reconstructed cluster of calorimeter energies associated with particles produced in the hadronization and decay of a bottom or charm quark.
 - [36] A b -tagged jet is one identified as consistent with that expected from the decay products of a bottom quark based on properties such as the presence of displaced track vertices or soft leptons.
 - [37] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005); A. Abulencia *et al.* (CDF Collaboration), *J. Phys. G* **34**, 2457 (2007).
 - [38] V.M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006); M. Abolins *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **584**, 75 (2008); R. Angstadt *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **622**, 298 (2010).
 - [39] V.M. Abazov *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **620**, 490 (2010).
 - [40] J. Freeman *et al.*, [arXiv:1205.1812](#); D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 052003 (2005); A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. D* **74**, 072006 (2006).

-
- [41] T. Aaltonen *et al.* (CDF and D0 Collaborations), [Phys. Rev. Lett.](#) **104**, 061802 (2010).
- [42] K. Nakamura *et al.* (Particle Data Group), [J. Phys. G](#) **37**, 075021 (2010); A. L. Read, [J. Phys. G](#) **28**, 2693 (2002); W. Fisher, Report No. FERMILAB-TM-2386-E.
- [43] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi, and M. Spira, [Eur. Phys. J. C](#) **71**, 1753 (2011).
- [44] S. Klimenko, J. Konigsberg, and T.M. Liss, Report No. FERMILAB-FN-0741 2003.
- [45] T. Andeen *et al.*, Report No. FERMILAB-TM-2365 2007.
- [46] L. Lyons, [Annals Applied Statistics](#) **2**, 887 (2008).
- [47] O.J. Dunn, [J. Am. Stat. Assoc.](#) **56**, 52 (1961).